

Short stature is associated with coronary heart disease: a systematic review of the literature and a meta-analysis

Tuula A. Paajanen^{1*}, Niku K.J. Oksala^{1,2}, Pekka Kuukasjärvi¹, and Pekka J. Karhunen¹

¹Department of Forensic Medicine, Medical School, University of Tampere and Centre for Laboratory Medicine, Tampere University Hospital, Tampere, Finland; and ²Division of Vascular Surgery, Department of Surgery, Tampere University Hospital, Tampere, Finland

Received 27 September 2009; revised 25 February 2010; accepted 23 March 2010; online publish-ahead-of-print 8 June 2010

See page 1674 for the editorial comment on this article (doi:10.1093/eurheartj/ehq183)

Aims

The aim of this study was to assess the relationship between short stature and coronary heart disease (CHD) morbidity and mortality.

Methods and results

We performed a systematic search from MEDLINE, PREMEDLINE, and All EBM Reviews as well as from a reference list of relevant articles. We used SPICO (Study design, Patient, Intervention, Control-intervention, Outcome) criteria. The methodological quality of studies was analysed by modified Borghoust criteria. From a total of 1907 articles, we selected 52 studies comprising population-based follow-up studies and patient cohorts followed after a CHD event, as well as case-control studies with height either as a continuous or categorical variable, totalling 3 012 747 individuals. The short ones were below 160.5 cm and tall ones over 173.9 cm on average. Among the shortest height category, the relative risks were 1.35 (95% CI 1.25–1.44) for all-cause mortality, 1.55 (1.37–1.74) for all cardiovascular disease (CVD) mortality, 1.49 (1.33–1.67) for CHD, and 1.52 (1.28–1.81) for myocardial infarction when compared with those within the highest height category. The mean relative risk was 1.46 (1.37–1.55). Short stature was associated with increased cardiovascular morbidity and mortality in both genders.

Conclusion

The relationship between short stature and CVD appears to be a real one. On the basis of comparison, adults within the shortest category had an ~50% higher risk of CHD morbidity and mortality than tall individuals.

Keywords

Short stature • Height • Coronary heart disease • Systematic review • Meta-analysis

Introduction

History

The first report on the inverse association between coronary heart disease (CHD) and height was published in 1951.¹ It was found that the average height of males hospitalized for myocardial infarction (MI) prior to the age of 40 was 5.08 cm lower (170.2 cm) than the average height of the non-hospitalized control group (175.3 cm). Since then, the association between short stature and cardiovascular diseases (CVDs) has been dealt with in more than 1900 papers according to database search. Many reports agree with Gertler *et al.*,^{2–12} but this association was also considered a misconception, as several large epidemiologic follow-up studies^{3,13–15} showed no association between height and CHD risk. It has even been suggested^{16,17} that racial factors may

determine the outcome, e.g. the shorter populations such as southern Europeans have lower death rates from CHD and all causes compared with, for example, the taller northern Europeans. Although seven reviews have been published,^{16–22} no systematic meta-analysis has been done on this topic.

The reasons for a systematic review

In this paper, we have performed, for the first time, a systematic review of studies reporting on the association between adult height and CHD events. What makes it difficult to investigate this association is that the literature deals with various populations and outcomes as well as different height criteria within various study designs. Some studies start with healthy people and follow their outcome in relation to their height, whereas other studies follow patients by stature after a CHD event. Thirdly, there are

* Corresponding author. Tel: +358 50 321 0023, Fax: +358 3 3551 7511, Email: tuula.paajanen@uta.fi

Published on behalf of the European Society of Cardiology. All rights reserved. © The Author 2010. For permissions please email: journals.permissions@oxfordjournals.org

case-control studies that comprise coronary artery disease patients and healthy controls defined by height, with varying outcomes and treatments. These studies also differ in the way they take the effect of known CHD risk factors into account, such as age, sex, smoking, lipid disorders, and diabetes—or whether they include any of these as confounding factors in their analyses. This is important, since in many papers the association between height and CHD often disappears when adjusted for all other factors. The most striking obstacle was, however, the use of various height categories.

Article quality was assessed by Quality of Reporting of Meta-analysis criteria. The hypothesis of this study was that short height is associated with increased CHD risk. We decided to compare the shortest group to the tallest group instead of using a fixed height limit and calculated average heights for combined risk values from the meta-analysis.

Objectives

This systematic review addresses whether short height shows an inverse association with variable CHD endpoints, and what the magnitude of the possible risk is. The meta-analysis will also address the effect of gender on the risk of different endpoints.

Criteria for considering studies for this review

Types of studies, participants, interventions, and outcome measures

According to the SPICO (Study design, Patient, Intervention, Control-intervention, Outcome) criteria, the original *study design* had to be a systematic review, meta-analysis, randomized clinical trial, clinical trial, and cohort or case-control setting and feature a number of subjects over 200 in total. *Patients* had to be either healthy at the beginning or already having symptomatic CHD. The patient setting was considered quite helpful if there was division into men vs. women. *Intervention and control intervention* had to be continuous, with the mean height and standard deviation given to be dichotomized as short vs. tall ones or classified and, if classified, there had to be at least two classes. *Outcome* had to be defined as diagnosis of angina pectoris, ischaemic heart disease (IHD) or heart disease without MI, acute MI, or history of MI, coronary artery occlusion equal to or more than 50%, revascularization or percutaneous transluminal coronary angioplasty (PTCA), as well as all-cause mortality, CVD mortality, or CHD mortality, or in clinical trials and cohort studies, any of the previous after at least 2 years of follow-up.

An article was excluded if the previously mentioned SPICO criteria were not fulfilled, or the information was insufficient for quality assessment or further conclusions, or height was only mentioned as a confounding factor.

Search strategy for identification of studies

We aimed to identify all articles that investigated height and CHD association by performing a systematic literature search from MEDLINE, PreMEDLINE, and All EBM Reviews, search ending 6 April 2007. The detailed search strategy is included as a Supplementary material online. This systematic search from the

sources above yielded a total of 1907 articles. On 3 December 2007, we checked that there were no new articles to include.

Two reviewer authors (T.A.P. and N.K.J.O.) independently assessed the studies for eligibility without consideration of the results. In cases of distributed opinions, a third evaluator was used (P.J.K.).

We included an article if predefined SPICO criteria were fulfilled in title and abstract. A total of 68 articles were reviewed as full-text versions. Next, we excluded articles that did not report adult height but birth height (five rejected on this ground) and articles that did not use English language (three rejected on this ground). Sixty articles remained after these additional restrictions.

After the database search, we screened the reference lists of included articles, which yielded 22 new articles. Excluding reviews and overlapping cohorts at this point, the number of eligible articles narrowed from 82 to 52, from which the final synthesis was made.

These articles comprised various racial and socioeconomic groups and samples from both genders separately totalling 3 012 747 individuals comprising population-based follow-up studies of initially healthy people, patient cohorts followed after a CHD event, and case-control studies with adequately defined classification of height either as a continuous or categorical variable. *Figure 1* summarizes the systematic forwarding of the search.

Methods

One author (T.A.P) extracted data from a previously designed Excel database see (Supplementary material online, *Table S1*). Completed data forms were checked for discrepancies by two authors (N.K.J.O. and P.J.K.).

Systematic review

We decided to do subgroup analyses by first extracting the data of those articles that reported risk ratios (RRs). If RRs were reported for the tallest group, we changed them to represent RRs for the shortest group. If RRs were not reported, but cross-tabulation for RR calculation was possible, we used the tallest group as the reference group and calculated RRs for the shortest group.

When odds ratio (OR) was reported for the tallest group, we used the following calculation: $RR = OR/[1 - p_c(1 - OR)]$, in which p_c is the event rate for the control group.²³ When possible, we extracted the gender-specific data because gender is one known confounding factor.

We also used ORs, hazard ratios (HRs), and other variables separately when it was impossible to calculate RRs due to the total number of individuals lacking. Thus, all 52 articles included are also evaluated in this systematic review results.

Meta-analysis

Meta-analysis was accomplished using statistical StatsDirect software. To examine the heterogeneity of the results and summarize results across various articles, we used methods suggested by DerSimonian and Laird.^{24–26} The I^2 test showed 72.1% inconsistency (95% CI 62.8–78.2%). Though we found significant within-trials and between-trials heterogeneity for the articles included in the meta-analysis, we present summary estimates based on a random effects model instead of a fixed effect model. We did not remove any of the selected studies based on heterogeneity. The Z-test for overall effect was 12.1 with $P < 0.0001$. According to bias analysis (*Figure 2*), our analysis was




Figure 1 The Quality of Reporting of Meta-analysis flow diagram.




Figure 2 Bias test.

unbiased (Begg–Mazumdar: Kendall's $\tau = 0.147449$, $P = 0.1175$ and Egger: bias = 0.724877 (95% CI = -0.172388–1.622143), $P = 0.111$).

Description of studies

Supplementary material online, *Table S1*, presents the most important descriptive information for the 52 studies included in our analysis. The total number of participants included in our systematic review was 3 012 747. In the meta-analysis, it was possible to include 22 studies see (Supplementary material online, *Table S2*). From a total of 30 studies, it was not possible to extract RRs and they were analysed in the text see (Supplementary material online, *Table S3*).

Methodological quality of included studies

Criteria for the assessment of the quality of studies were modified from Borghoust *et al.*²⁷ see (Supplementary material online, *Table S4*). The quality of the studies included in the systematic review varied from 7 to 15 see (Supplementary material online, *Table S1*). The quality score was commonly low because the subcriteria of the study size (5c, 5b) or the amount of dropouts/loss to follow-up (8a) were too small, or information provided from dropouts/loss to follow-up (8b) was insufficient. There was no association between quality score and bias assessment plot scattering (data not shown).

Results

Studies included in meta-analysis

Of the 52 studies, 22^{4,5,7,11,28–45} provided adequate data for a meta-analysis based on RRs see (*Figure 3*, Supplementary material online, *Table S2*). Meta-analysis shows that risks for different kinds of cardiovascular endpoints are higher within the shortest ones when compared with the tallest height categories. The combined RR was 1.46 (95% CI 1.37–1.55). On average, the short ones were below 160.5 cm and tall ones over 173.9 cm, so the average height cut-off for short ones was 13.4 cm lower than the average height cut-off for the tall ones. When men and women were considered independently on average, short men were below 165.4 cm and short women below 153.0 cm, tall men over 177.5 cm and tall women over 166.4 cm. Height cut-offs were available for 21 articles. Only one article³³ failed to report height

cut-offs and was not taken into the calculations. Additional meta-analysis information is provided in Supplementary material online, *Table S2*.

The combined RR for all-cause mortality for short men was 1.37 (1.29–1.46) and for short women 1.55 (1.41–1.70) see (Supplementary material online, *Figures S1* and *S2*). On average, the short ones were defined to be below 161.1 cm and the tall ones over 176.0 cm and short men below 165.5 cm and short women below 153.3 cm, tall men over 178.9 cm and tall women over 163.7 cm see (Supplementary material online, *Figure S3*).

Similarly, the combined risk for all types of cardiovascular (CVD) deaths among men and women was 1.55 (95% CI 1.37–1.74) see (Supplementary material online, *Figure S4*). In line, the risk for combined CHD mortality and morbidity brought by short stature among men was 1.49 (1.33–1.67), varying between 1.10 and 4.55. In these studies, men with a height of <166.1 cm were considered short and men >176.2 cm tall. There were only three studies in women reporting a risk varying from 1.10 to 2.08 for short ones (<153.1 cm) compared with the risk among tall ones (>165.6 cm) see (Supplementary material online, *Figure S5*).

The risk of MI incidence associated with short stature was 1.52 (95% CI 1.28–1.81) for all. In these studies, the mean reported height cut-off for short males was 14.4 cm lower (164.0 cm) than the average height cut-off of tall men (178.4 cm). Similarly, there was a 14.5 cm difference between reported cut-offs for short (153.7 cm) and tall women (168.2 cm) see (Supplementary material online, *Figure S7*).

Meta-analysis subgroup results see (Supplementary material online, *Figures 1–8*) show a more detailed interpretation.

Studies not included in the meta-analysis

There were 30 studies fulfilling the SPICO criteria but not included in the meta-analysis^{2,3,6,8–10,12–15,46–65} because it was impossible to calculate RRs for absolute height cut-offs see (Supplementary material online, *Table S3*). Of these, seven studies reported RRs by variable increments of height without data on measured height.^{8,13,15,47,51,61,64} From these seven studies, we can conclude that a 5–10 cm height increase is usually linked with a reduction




Figure 3 Meta-analysis of 22 articles.

of RR for all-cause death or CHD by 10–15% and a 30 cm decrease of height may double the risk.

Additionally, six studies reported ORs from which RRs could not be calculated.^{3,50,52,53,56,65} Incidence of non-fatal MI and CHD as well as coronary lesions $\geq 50\%$ were inversely associated with height, but 30-day or operative mortality after isolated coronary artery bypass grafting did not show significant association with height.

Furthermore, three studies reported HRs by quartiles^{14,60,63} and as with ORs, some researchers also used staircase increments of height when reporting HRs.^{6,49,55,59} An inverse association for CHD was found when height was used as a continuous variable in every study; in a twin study, discordant for height was found not only between twin individuals but also within twin pairs. Per 5 or 10 or 15 cm increase HR was inversely associated with height for fatal CVD and for fatal CHD and for all-cause mortality.

There were four studies^{2,9,10,48} reporting only statistically significant *P*-values to support statements of height being an important factor for risk of CHD.

Six studies could not be grouped at all, based on the used endpoints.^{12,46,54,57,58,62} However, reported endpoints such as operative mortality percentage after coronary artery bypass grafting (CABG), advanced coronary lesions in an autopsy study, incidence of IHD, estimated CHD mortality ratio, all-cause mortality ratio, and affirmative answer about symptoms implying coronary or peripheral atherosclerotic disease were all inversely associated with height in these studies.

Discussion

In this paper, we have addressed two questions: Is adult short stature really a risk factor for CHD events and, if it is, how big a factor is it? Our systematic review and meta-analysis show that adult short stature poses ~ 1.5 times higher risk for CHD morbidity and mortality than being a tall individual. This appears to be true both for men and women and for different kinds of endpoints. Although it has been previously hypothesized that height may possess a greater risk for men, shorter women may pose an even higher risk when other confounding factors are not taken into consideration.

Due to the heterogeneity of studies, we cannot reliably answer the question on the critical absolute height. The height cut-off points did not only differ between the articles but also between men and women and between ethnic groups. This is why we used the shortest-vs.-tallest group setting. Our strategy thereby reduced the study population pool. Based on the bias analysis, we used the random effect model, assuming that the true height value varies in different study populations. The advantage of the use of random effect model was that we did not have to remove any of the selected studies based on the variability mentioned above. We believe that this heterogeneity only strengthens the association found.

According to our modified Borghoust criteria list, the methodological quality of the articles appeared to be reasonably good on average to be used for the assessment of height as a risk factor, though most of them were originally designed for other purposes. Two of the most common methodological shortcomings appeared to be the missing number of individuals with outcome and lack of information provided for dropouts/loss to follow-up. More than half the included studies failed to fulfil these criteria.

Our systematic review has some limitations. Meta-analyses are also prone to different kinds of biases and confounding factors that are inherent to the original studies. The possibility of missing published studies also remains, though we used a large set of keywords for databases and standardized the search flow in accordance with the guidelines. Furthermore, we searched only for studies which were published in indexed journals or found from reference lists, so unpublished studies and non-indexed journals may have been partly missed. Positive results are more easily published, but in this systematic review, height was usually not the main topic of the original paper but rather was given as a demographic variable. This is why we believe that our results were not biased by positive publication bias. Only a few studies were focused to identify height as a prognostic factor.

We excluded only three studies out of the total of 1907 articles because they were not published in English. This language limit would hardly have changed the result because in these abstracts the results were parallel.

The reason(s) for short stature being a risk factor for CHD remain open for hypotheses. Most commonly, in previous studies it has been suggested that the reason behind this association could be low socioeconomic background with associated risk factors such as poor nutrition and infections resulting in poor foetal or early-life growth.^{39,66}

Nwasokva *et al.*⁵⁶ found that short men have not only a higher prevalence of coronary disease, but also a greater severity of coronary disease than tall men.⁵⁶ Also, Kortelainen and Särkioja¹² found that short stature was associated with more advanced coronary lesions.

In Physicians Health Study, there was a significant association between height and MI, but not between height and cardiovascular death.⁶⁷ Similarly, the Framingham Heart Study found no association between short stature and increased risk for all-cause or cardiovascular mortality in either sex, although shorter women had increased risk for MI.³³

In the Coronary Artery Surgery Study (CASS), surgical mortality was inversely related to the average diameter of the grafted coronary arteries in both men and women.⁶⁸ It was therefore hypothesized that the physical size of the patient, including coronary artery diameter, may predict operative mortality.⁶⁸ In recent studies using angiographic measurements, the coronary artery diameter was correlated with height and body weight.^{69,70} It could be hypothesized that smaller coronary arteries may be occluded earlier in life under similar risk conditions.

Conclusions

The relationship between short stature and CVD seems to be a real one. Adults within the shortest category had a $\sim 50\%$ higher risk of CHD morbidity and mortality compared with tall individuals. The possible pathophysiological, environmental, and genetic background of this peculiar association is not known.

Implications for practice

Height is used to calculate body mass index (BMI), which is a widely used quantity risk of CHD. The value of BMI has been recently questioned by reports showing that BMI may not associate

with the severity of CHD in angina patients with chronic kidney disease⁷¹ or may even be inversely associated.^{72,73} This has been discussed as the obesity paradox. The results of this meta-analysis suggest that height may be considered as a possible independent factor to be used in CHD risk calculations.

Implications for research

It would be interesting to explore—e.g. in autopsy series—the possibility that short stature is connected with the risk of CHD and MI through the effect of smaller coronary artery diameter, and that smaller coronary arteries may be occluded earlier in life under similar risk conditions. Recent findings on the genetic background of body height⁷⁴ suggest that inherited factors rather than speculative early-life poor nutrition or birth weight may explain the association between small stature and later-life increased risk for CHD events.

Contributions

T.A.P., N.K.J.O., P.J.K., and P.K. designed the present study. T.A.P. analysed and interpreted the data, and drafted the article. N.K.J.O., P.J.K., and P.K. interpreted and organized the data, critically revised the article and contributed to the final version.

Supplementary material

Supplementary material is available at *European Heart Journal* online.

Funding

This study was supported by grants from the Pirkanmaa Regional Fund of the Finnish Cultural Foundation, the Medical Research Fund of Tampere University Hospital, the Aarno Koskelo Foundation, and the Finnish Foundation for Cardiovascular Research.

Conflicts of interest: none declared.

References

- Gertler MM, Garn SM, White PD. Young candidates for coronary heart disease. *JAMA* 1951;**147**:621–625.
- Blümchen G, Jetté M. Relationship between stature and coronary heart disease in a German male population. *Int J Cardiol* 1992;**36**:351–355.
- Christakis GT, Weisel RD, Buth KJ, Fremes SE, Rao V, Panagiotopoulos KP, Ivanov J, Goldman BS, David TE. Is body size the cause for poor outcomes of coronary artery bypass operations in women? *J Thorac Cardiovasc Surg* 1995;**110**:1344–1356. discussion 1356–1358.
- D'Avanzo B, La Vecchia C, Negri E. Height and the risk of acute myocardial infarction in Italian women. *Soc Sci Med* 1994;**38**:193–196.
- DaveySmith G, Hart C, Upton M, Hole D, Gillis C, Watt G, Hawthorne V. Height and risk of death among men and women: aetiological implications of associations with cardiorespiratory disease and cancer mortality. *J Epidemiol Community Health* 2000;**54**:97–103.
- Ferrie JE, Langenberg C, Shipley MJ, Marmot MG. Birth weight, components of height and coronary heart disease: evidence from the Whitehall II study. *Int J Epidemiol* 2006;**35**:1532–1542.
- Forsen T, Eriksson J, Qiao Q, Tervahauta M, Nissinen A, Tuomilehto J. Short stature and coronary heart disease: a 35-year follow-up of the Finnish cohorts of The Seven Countries Study. *J Intern Med* 2000;**248**:326–332.
- Glynn RJ, Rosner B. Comparison of risk factors for the competing risks of coronary heart disease, stroke, and venous thromboembolism. *Am J Epidemiol* 2005;**162**:975–982.
- Greig M, Pemberton J, Hay I, MacKenzie G. A prospective study of the development of coronary heart disease in a group of 1202 middle-aged men. *J Epidemiol Community Health* 1980;**34**:23–30.
- He GW, Acuff TE, Ryan WH, Bowman RT, Douthit MB, Mack MJ. Determinants of operative mortality in elderly patients undergoing coronary artery bypass grafting. Emphasis on the influence of internal mammary artery grafting on mortality and morbidity. *J Thorac Cardiovasc Surg* 1994;**108**:73–81.
- Hippe M, Vestbo J, Hein HO, Borch-Johnsen K, Jensen G, Sorensen TI. Familial predisposition and susceptibility to the effect of other risk factors for myocardial infarction. *J Epidemiol Community Health* 1999;**53**:269–276.
- Kortelainen ML, Särkioja T. Coronary atherosclerosis associated with body structure and obesity in 599 women aged between 15 and 50 years. *Int J Obes Relat Metab Disord* 1999;**23**:838–844.
- Krahn AD, Manfreda J, Tate RB, Mathewson FAL, Cuddy TE. Evidence that height is an independent risk factor for coronary artery disease (the Manitoba Follow-Up Study). *Am J Cardiol* 1994;**74**:398–399.
- Ness AR, Gunnell D, Hughes J, Elwood PC, Smith GD, Burr ML. Height, body mass index, and survival in men with coronary disease: follow up of the diet and reinfarction trial (DART). *J Epidemiol Community Health* 2002;**56**:218–219.
- Rosenberg CR, Shore RE, Pasternack BS. Height and mortality after myocardial infarction. *J Community Health* 1995;**20**:335–343.
- Samaras TT, Elrich H. Height, body size and longevity. *Acta Med Okayama* 1999;**53**:149–169.
- Samaras TT, Harold E, Storms LH. Is short height really a risk factor for coronary heart disease and stroke mortality? A review. *Med Sci Monit* 2004;**10**:RA63–RA76.
- Elford J, Whincup P, Shaper AG. Early life experience and adult cardiovascular disease: longitudinal and case-control studies. *Int J Epidemiol* 1991;**20**:833–844.
- Samaras TT. How body height and weight affect our performance, longevity, and survival. *J Wash Acad Sci* 1996;**84**:131–156.
- Samaras TT. Height, body size, and longevity: is smaller better for the human body? *West J Med* 2002;**176**:206–208.
- Samaras TT, Harold E, Storms LH. Is height related to longevity? *Life Sci* 2003;**72**:1781–1802.
- Williams SRP, Jones E, Bell W, Davies B, Bourne MW. Body habitus and coronary heart disease in men. A review with reference to methods of body habitus assessment. *Eur Heart J* 1997;**18**:376–393.
- Deeks JJ, Altman DG. Effect measures for meta-analysis of trials with binary outcomes. In: Egger M, Smith GD, Altman DG, eds. *Systematic Reviews in Health Care Meta-analysis in Context*. 2nd ed. London: BMJ Books; 2001. p313–p335.
- DerSimonian R, Laird N. Meta-analysis in clinical trials. *Control Clin Trials* 1986;**7**:177–188.
- Thompson SG. Why and how sources of heterogeneity should be investigated. In: Egger M, Smith GD, Altman DG, eds. *Systematic Reviews in Health Care Meta-analysis in Context*. 2nd ed. London: BMJ Books; 2001. p157–p175.
- Sterne JAC, Egger M, Smith GD. Investigating and dealing with publication and other biases. In: Egger M, Smith GD, Altman DG, eds. *Systematic Reviews in Health Care Meta-analysis in Context*. 2nd ed. London: BMJ Books; 2001. p189–p208.
- Borghoust JA, Koes BW, Bouter LM. The clinical course and prognostic factors of non-specific neck pain: a systematic review. *Pain* 1998;**77**:1–13.
- Allebeck P, Bergh C. Height, body mass index and mortality: do social factors explain the association? *Public Health* 1992;**106**:375–382.
- Cook NR, Hebert PR, Satterfield S, Taylor JO, Buring JE, Hennekens CH. Height, lung function and mortality from cardiovascular disease among the elderly. *Am J Epidemiol* 1994;**139**:1066–1076.
- Goldbourt U, Tanne D. Body height is associated with decreased long-term stroke but not coronary heart disease mortality? *Stroke* 2002;**33**:743–748.
- Helmert U, Shea S. Relation between body height and self-reported myocardial infarction in Germany. *Rev Environ Health* 1997;**12**:125–130.
- IONA Study Group. Determinants of coronary events in patients with stable angina: results from the impact of nicorandil in angina study. *Am Heart J* 2005;**150**:689.
- Kannam JP, Levy D, Larson M, Wilson PW. Short stature and risk for mortality and cardiovascular disease events. The Framingham Heart Study [see comment]. *Circulation* 1994;**90**:2241–2247.
- LaVecchia C, Decarli A, Negri E, Ferraroni M, Pagano R. Height and the prevalence of chronic disease. *Rev Epidém et Santé Publ* 1992;**40**:6–14.
- McCarron P, Okasha M, McEwen J, Smith GD. Height in young adulthood and risk of death from cardiorespiratory disease: a prospective study of male former students of Glasgow University, Scotland [see comment]. *Am J Epidemiol* 2002;**155**:683–687.
- Mukamal KJ, Maclure M, Sherwood JB, Kannam JP, Muller JE, Mittleman MA. Height is not associated with long-term survival after acute myocardial infarction. *Am Heart J* 2001;**142**:852–856.
- Myers WO, Davis K, Foster ED, Maynard C, Kaiser GC. Surgical survival in the Coronary Artery Surgery Study (CASS) registry. *Ann Thorac Surg* 1985;**40**:245–260.

38. Osika W, Ehlin A, Montgomery SM. Does height modify the risk of angina associated with economic adversity? *Econ Hum Biol* 2006;**4**:398–411.
39. Palmer JR, Rosenberg L, Shapiro S. Stature and the risk of myocardial infarction in women [see comment]. *Am J Epidemiol* 1990;**132**:27–32.
40. Parker DR, Lapane KL, Lasater TM, Carleton RA. Short stature and cardiovascular disease among men and women from two southeastern New England communities. *Int J Epidemiol* 1998;**27**:970–975.
41. Rich-Edwards JW, Manson JE, Stampfer MJ, Colditz GA, Willett WC, Rosner B, Speizer FE, Hennekens CH. Height and the risk of cardiovascular disease in women [see comment]. *Am J Epidemiol* 1995;**142**:909–917.
42. Rimm EB, Stampfer MJ, Giovannucci E, Ascherio A, Spiegelman D, Colditz GA, Willett WC. Body size and fat distribution as predictors of coronary heart disease among middle-aged and older US men. *Am J Epidemiol* 1995;**141**:1117–1127.
43. Song YM, Smith GD, Sung J. Adult height and cause-specific mortality: a large prospective study of South Korean men. *Am J Epidemiol* 2003;**158**:479–485.
44. Weintraub WS, Wenger NK, Kosinski AS, Douglas JS, Liberman HA, Morris DC, King SB III. Percutaneous transluminal coronary angioplasty in women compared with men. *J Am Coll Cardiol* 1994;**24**:81–90.
45. Yarnell JWG, Limb ES, Layzell JM, Baker IA. Height: a risk marker for ischaemic heart disease: prospective results from the Caerphilly and Speedwell Heart Disease Studies. *Eur Heart J* 1992;**13**:1602–1605.
46. Grover FL, Johnson RR, Marshall G, Hammermeister KE, Department of Veterans Affairs Cardiac Surgeons. Factors predictive of operative mortality among coronary artery bypass subsets. *Ann Thorac Surg* 1993;**56**:1296–1307.
47. Jousilahti P, Tuomilehto J, Vartiainen E, Eriksson J, Puska P. Relation of adult height to house-specific and total mortality: a prospective follow-up study of 31,199 middle-aged men and women in Finland. *Am J Epidemiol* 2000;**151**:1112–1120.
48. Kelly RF, Mohanty J, Hashim AS, Parrillo JE. Association between height and coronary artery disease in black men and women. *Am J Cardiol* 2000;**85**:1253–1255.
49. Langenberg C, Shipley MJ, Batty GD, Marmot MG. Adult socioeconomic position and the association between height and coronary heart disease mortality: findings from 33 years of follow-up in the Whitehall Study. *Am J Public Health* 2005;**95**:628–632.
50. Lawlor DA, Taylor M, Smith GD, Gunnell D, Ebrahim S. Associations of components of adult height with coronary heart disease in postmenopausal women: the British women's heart and health study. *Heart* 2004;**90**:745–749.
51. Liao Y, McGee DL, Cao G, Cooper RS. Short stature and risk of mortality and cardiovascular disease: negative findings from the NHANES I epidemiologic follow-up study. *J Am Coll Cardiol* 1996;**27**:678–682.
52. Lundberg M, Diderichsen F, Hallqvist J for the SHEEP Study Group. Is the association between short stature and myocardial infarction explained by childhood exposures—a population-based case referent study (SHEEP). *Scand J Public Health* 2002;**30**:249–258.
53. Mehta RH, Califf RM, Garg J, White HD, de Werf FV, Armstrong PW, Pieper KS, Topol EJ, Granger CB. The impact of anthropomorphic indices on clinical outcomes in patients with acute ST-elevation myocardial infarction. *Eur Heart J* 2007;**28**:415–424.
54. Morris JN, Kagan A, Pattison DC, Gardner MJ, Raffle PAB. Incidence and prediction of ischaemic heart-disease in London busmen. *Lancet* 1966;**2**:553–559.
55. Morris RW, Whincup PH, Emberson JR, Lampe FC, Walker M, Shaper AG. North-South gradients in Britain for stroke and CHD: are they explained by the same factors? *Stroke* 2003;**34**:2604–2609.
56. Nwasokva ON, Weiss M, Gladstone C, Bodenheimer MM. Higher prevalence and greater severity of coronary disease in short versus tall men referred for coronary arteriography. *Am Heart J* 1997;**133**:147–152.
57. Paffenbarger RS, Wing AL. Chronic disease in former college students: X the effects of single and multiple characteristics on risk of fatal coronary heart disease. *Am J Epidemiol* 1969;**90**:527–535.
58. Peck AM, Vagero DH. Adult body height, self perceived health and mortality in the Swedish population. *J Epidemiol Community Health* 1989;**43**:380–384.
59. Pocock SJ, McCormack V, Gueyffier F, Boutitie F, Fagard RH, Boissel JP on behalf of the INDANA project steering committee. A score for predicting risk of death from cardiovascular disease in adults with raised blood pressure, based on individual patient data from randomised controlled trials [see comment]. *BMJ* 2001;**323**:75–81.
60. Silventoinen K, Zdravkovic S, Skytthe A, McCarron P, Herskind AM, Koskenvuo M, de Faire U, Pedersen N, Christensen K, Kaprio J. Association between height and coronary heart disease mortality: a prospective study of 35,000 twin pairs. *Am J Epidemiol* 2006;**163**:615–621.
61. Strandberg TE. Inverse relation between height and cardiovascular mortality in men during 30-year follow-up. *Am J Cardiol* 1997;**80**:349–350.
62. Thelle DS, Førde OH, Try K, Lehmann EH. The Tromsø heart study. Methods and main results of the cross-sectional study. *Acta Med Scand* 1976;**200**:107–118.
63. Tunstall-Pedoe H, Woodward M, Tavendale R, A'Brook R, McCluskey MK. Comparison of the prediction by 27 different factors of coronary heart disease and death in men and women of the Scottish Heart Health Study: cohort study [erratum appears in *BMJ* 1998 Jun 20;316(7148):1881]. *BMJ* 1997;**315**:722–729.
64. Waaler HT. Height, weight and mortality. The Norwegian experience. *Acta Med Scand* 1984;(Suppl. 679):3–56.
65. Wamala SP, Lynch J, Kaplan GA. Women's exposure to early and later life socioeconomic disadvantage and coronary heart disease risk: the Stockholm Female Coronary Risk Study. *Int J Epidemiol* 2001;**30**:275–284.
66. Barker DJP. The intrauterine origins of cardiovascular and obstructive lung disease in adult life. *J R Coll Phys Lond* 1991;**25**:129–133.
67. Hebert PR, Rich-Edwards JW, Manson JE, Ridker PM, Cook NR, ÓConnor GT, Buring JE, Hennekens CH. Height and incidence of cardiovascular disease in male physicians. *Circulation* 1993;**88**:1437–1443.
68. Fisher LD, Kennedy JW, Davis KB, Maynard C, Fritz JK, Kaiser G, Myers WO, and the Participating CASS Clinics. Association of sex, physical size, and operative mortality after coronary artery bypass in the Coronary Artery Surgery Study (CASS). *J Thorac Cardiovasc Surg* 1982;**84**:334–341.
69. Lemos PA, Ribeiro EE, Perin MA, Kajita LJ, de Magalhães MA, Falcão JLAA, Filho AE, da Gama MN, Horta PE, Marchiori GG, Spadaro AG, Martinez EE. Angiographic segment size in patients referred for coronary intervention is influenced by constitutional, anatomical, and clinical features. *Int J Cardiovasc Imaging* 2007;**23**:1–7.
70. Zeina AR, Rosenschein U, Barmer E. Dimensions and anatomic variations of left main coronary artery in normal population: multidetector computed tomography assessment. *Coron Artery Dis* 2007;**18**:477–482.
71. Wu IW, Hung MJ, Chen YC, Cheng WJ, Wu MS. High body mass index is not associated with coronary artery disease in angina patients with chronic kidney disease: a coronary angiography study. *Am J Med Sci* 2008;**336**:303–308.
72. Venkatesh PK, Molnar J, Swaminathan PD, Khosla S, Arora RR. Normal coronary arteries in patients with systolic heart failure who have higher body mass index. *Heart Lung* 2007;**36**:125–131.
73. Rubinshtein R, Halon DA, Jaffe R, Shahla J, Lewis BS. Relation between obesity and severity of coronary artery disease in patients undergoing coronary angiography. *Am J Cardiol* 2006;**97**:1277–1280.
74. Sammalisto S, Hiekkalinna T, Schwander K, Kardia S, Weder AB, Rodriguez BL, Doria A, Kelly JA, Bruner GR, Harley JB, Redline S, Larkin EK, Patel SR, Ewan AJH, Weber JL, Perola M, Peltonen L. Genome-wide linkage screen for stature and body mass index in 3,032 families: evidence for sex- and population-specific genetic effects. *Eur J Hum Genet* 2008;**17**:258–266.